

No part of this product may be reproduced in any form or by any electronic or mechanical means, including information storage and retrieval systems, without written permission from the IB.

Additionally, the license tied with this product prohibits commercial use of any selected files or extracts from this product. Use by third parties, including but not limited to publishers, private teachers, tutoring or study services, preparatory schools, vendors operating curriculum mapping services or teacher resource digital platforms and app developers, is not permitted and is subject to the IB's prior written consent via a license. More information on how to request a license can be obtained from https://ibo.org/become-an-ib-school/ib-publishing/licensing/applying-for-a-license/.

Aucune partie de ce produit ne peut être reproduite sous quelque forme ni par quelque moyen que ce soit, électronique ou mécanique, y compris des systèmes de stockage et de récupération d'informations, sans l'autorisation écrite de l'IB.

De plus, la licence associée à ce produit interdit toute utilisation commerciale de tout fichier ou extrait sélectionné dans ce produit. L'utilisation par des tiers, y compris, sans toutefois s'y limiter, des éditeurs, des professeurs particuliers, des services de tutorat ou d'aide aux études, des établissements de préparation à l'enseignement supérieur, des fournisseurs de services de planification des programmes d'études, des gestionnaires de plateformes pédagogiques en ligne, et des développeurs d'applications, n'est pas autorisée et est soumise au consentement écrit préalable de l'IB par l'intermédiaire d'une licence. Pour plus d'informations sur la procédure à suivre pour demander une licence, rendez-vous à l'adresse suivante : https://ibo.org/become-an-ib-school/ib-publishing/licensing/applying-for-a-license/.

No se podrá reproducir ninguna parte de este producto de ninguna forma ni por ningún medio electrónico o mecánico, incluidos los sistemas de almacenamiento y recuperación de información, sin que medie la autorización escrita del IB.

Además, la licencia vinculada a este producto prohíbe el uso con fines comerciales de todo archivo o fragmento seleccionado de este producto. El uso por parte de terceros —lo que incluye, a título enunciativo, editoriales, profesores particulares, servicios de apoyo académico o ayuda para el estudio, colegios preparatorios, desarrolladores de aplicaciones y entidades que presten servicios de planificación curricular u ofrezcan recursos para docentes mediante plataformas digitales— no está permitido y estará sujeto al otorgamiento previo de una licencia escrita por parte del IB. En este enlace encontrará más información sobre cómo solicitar una licencia: https://ibo.org/become-an-ib-school/ib-publishing/licensing/applying-for-a-license/.



Chemistry Higher level Paper 2

Thursday 5 November 2020 (afternoon)

		Cai	lulua	ie se	55101	Hull	ibei	
ſ								
L								
-						•		

2 hours 15 minutes

Instructions to candidates

- Write your session number in the boxes above.
- Do not open this examination paper until instructed to do so.
- Answer all questions.
- Answers must be written within the answer boxes provided.
- A calculator is required for this paper.
- A clean copy of the **chemistry data booklet** is required for this paper.
- The maximum mark for this examination paper is [90 marks].

245004



-2- 8820-6102

Please do not write on this page.

Answers written on this page will not be marked.



24FP02


State the full electron configuration of the chlorine atom. State, giving a reason, whether the chlorine atom or the chloride ion has a larger radius.	[1]
	[1]
ii) Outline why the chlorine atom has a smaller atomic radius than the sulfur atom.	[2]
	i) Outline why the chlorine atom has a smaller atomic radius than the sulfur atom.



Turn over

(Question 1 continued)

(iv) The mass spectrum of chlorine is shown.



Outline the reason for the two peaks at m/z = 35 and 37.

[1]

.....

(v) Explain the presence and relative abundance of the peak at m/z = 74. [2]

(b) 2.67 g of manganese(IV) oxide was added to 200.0 cm³ of 2.00 mol dm⁻³ HCl.

 $MnO_2(s) + 4HCl(aq) \rightarrow Cl_2(g) + 2H_2O(l) + MnCl_2(aq)$

(i) Calculate the amount, in mol, of manganese(IV) oxide added. [1]

.....



(Question 1 continued) Determine the limiting reactant, showing your calculations. [2] (iii) Determine the excess amount, in mol, of the other reactant. [1] Calculate the volume of chlorine, in dm³, produced if the reaction is conducted at (iv) standard temperature and pressure (STP). Use section 2 of the data booklet. [1] State the oxidation state of manganese in $\mathrm{MnO_2}$ and $\mathrm{MnCl_2}$. (v) [2] MnO₂:

(vi) Deduce, referring to oxidation states, whether MnO₂ is an oxidizing or reducing agent. [1]

(This question continues on the following page)

MnCl₂:



Turn over

(Question 1 continued)

(c)	Chlo	rine gas reacts with water to produce hypochlorous acid and hydrochloric acid.	
		$Cl_2(g) + H_2O(l) \rightleftharpoons HClO(aq) + HCl(aq)$	
	(i)	Hypochlorous acid is considered a weak acid. Outline what is meant by the term weak acid.	[1]
	(ii)	State the formula of the conjugate base of hypochlorous acid.	[1]
	(iii)	Calculate the concentration of H^+ (aq) in a HClO(aq) solution with a pH = 3.61.	[1]
(d)	(i)	State the type of reaction occurring when ethane reacts with chlorine to produce chloroethane.	[1]
	(ii)	Predict, giving a reason, whether ethane or chloroethane is more reactive.	[1]



Ouestion	1	continued)	
Question		continued)	

(iii)	Explain the mechanism of the reaction between chloroethane and aqueous sodium hydroxide, NaOH (aq), using curly arrows to represent the movement of electron pairs.	[3]
(iv)	Ethoxyethane (diethyl ether) can be used as a solvent for this conversion. Draw the structural formula of ethoxyethane.	[1]
(v)	Deduce the number of signals and chemical shifts with splitting patterns in the ¹ H NMR spectrum of ethoxyethane. Use section 27 of the data booklet.	[3]



(Question 1 continued)

(e)	CCI	₂ F ₂ is a common chlorofluorocarbon, CFC.	
	(i)	Calculate the percentage by mass of chlorine in CCl ₂ F _{2.}	[2]
	(ii)	Comment on how international cooperation has contributed to the lowering of CFC emissions responsible for ozone depletion.	[1]
	(iii)	CFCs produce chlorine radicals. Write two successive propagation steps to show how chlorine radicals catalyse the depletion of ozone.	[2]



2. Compound A is in equilibrium with compound B.



(a) Predict the electron domain and molecular geometries around the **oxygen** atom of molecule **A** using VSEPR.

[2]

Electron domain geometry:	
Molecular geometry:	


- (b) State the type of hybridization shown by the central carbon atom in molecule **B**. [1]
- (c) State the number of sigma (σ) and pi (π) bonds around the central carbon atom in molecule **B**. [1]

σ-bonds:

 π -bonds:

(Question 2 continued)

(d) The IR spectrum of one of the compounds is shown:



Deduce, giving a reason, the compound producing this spectrum.

[1]

٠.	 •	 ٠.	•	 •	•	 •	•	 •	•	•	 •	•	•	 •	•	•	•	•	 •	•	•	 •	•	 	•	•		•	•	•	 •	•	•	•	 •	•	•	 	•	•	•	
	 -	 																						 			 											 				

(e) Compound **A** and **B** are isomers. Draw two other structural isomers with the formula C_3H_6O .

[2]



(Question 2 continued)

-			•	
	(f)	(i)	The equilibrium constant, K_c , for the conversion of A to B is 1.0×10^8 in water at 298 K.	
			Deduce, giving a reason, which compound, A or B , is present in greater concentration when equilibrium is reached.	[1]
		(ii)	Calculate the standard Gibbs free energy change, ΔG^{\ominus} , in kJ mol ⁻¹ , for the reaction (A to B) at 298 K. Use sections 1 and 2 of the data booklet.	[1]
	(g)	Prop	panone can be synthesized in two steps from propene.	
		(i)	Suggest the synthetic route including all the necessary reactants and steps.	[3]
I				



Turn over

	(ii) Suggest why propanal is a minor product obtained from the synthetic route in (g)(i).	[2]
3.	An equation for the combustion of propane is given below.	
	$C_3H_8(g) + 5O_2(g) \rightarrow 3CO_2(g) + 4H_2O(g)$	
	(a) Determine the standard enthalpy change, ΔH^{\ominus} , for this reaction, using section 11 of the data booklet.	[3]
	(b) Calculate the standard enthalpy change, ΔH^{\ominus} , for this reaction using section 12 of the data booklet.	[2]



(Question 3 continued)

(c)	Predict, giving a reason, whether the entropy change, ΔS^{\ominus} , for this reaction is negative or positive.	[1]
(d)	Calculate ΔS^{\ominus} for the reaction in J K ⁻¹ , using section 12 of the data booklet.	
	The standard molar entropy for oxygen gas is 205 J K ⁻¹ mol ⁻¹ .	[2]
(e)	Calculate the standard Gibbs free energy change, ΔG^{\ominus} , in kJ , for the reaction at 5°C, using your answers to (b) and (d). Use section 1 of the data booklet.	
	(If you did not obtain an answer to (b) or (d) use values of $-1952\mathrm{kJ}$ and $+113\mathrm{JK^{-1}}$ respectively, although these are not the correct answers.)	[2]



Turn over

4. Nickel catalyses the conversion of propanone to propan-2-ol.

0	H ₂	OH I
	Ni, heat	

(a)	Outline how a catalyst increases the rate of reaction.	[1]
(b)	Explain why an increase in temperature increases the rate of reaction.	[2]

and propan-2-ol.	[3]

Discuss, referring to intermolecular forces present, the relative volatility of propanone

.....

(This question continues on the following page)

(c)




(Question 4 continued)

The diagram shows an unlabelled voltaic cell for the reaction. (d)

$$Pb^{2+}(aq) + Ni(s) \rightarrow Ni^{2+}(aq) + Pb(s)$$

Label the diagram with the species in the equation.

[1]



(ii)	Calculate the standard cell potential, in V, for the cell at 298 K. Use section 24 of
	the data booklet.

[1]

(iii)	Calculate the standard free energy change, ΔG^{\ominus} , in kJ , for the cell using sections	
	1 and 2 of the data booklet.	[1

г	,	1
ı		ı
L		•

(iv)	Suggest a metal that could replace nickel in a new half-cell and reverse the
	electron flow. Use section 25 of the data booklet.

Г	•	1	
ı			
L			



Turn over


(Question	4	continu	ed)
, ~ ~ ~ ~			,

	(v)	Describe the bonding in metals.	[2]
	(vi)	Nickel alloys are used in aircraft gas turbines. Suggest a physical property altered by the addition of another metal to nickel.	[1]
5.		t performs a titration to determine the concentration of ethanoic acid, CH ₃ COOH, in using potassium hydroxide.	
	(a) Wr	te a balanced equation for the reaction.	[1]



(Question 5 continued)

(b) The pH curve for the reaction is given.



(i) Identify the **major** species, other than water and potassium ions, at these points. [2]

B:		
C:		

(ii) State a suitable indicator for this titration. Use section 22 of the data booklet. [1]

(iii) Suggest, giving a reason, which point on the curve is considered a buffer region. [1]

(This question continues on the following page)



Turn over

Question 5	continu	ed)
------------	---------	-----

(c)	State the K_a expression for ethanoic acid.	[1]
(d)	Calculate the $K_{\rm b}$ of the conjugate base of ethanoic acid using sections 2 and 21 of the data booklet.	[1]
(e)	In a titration, 25.00 cm³ of vinegar required 20.75 cm³ of 1.00 mol dm⁻³ potassium hydroxide to reach the end-point.	
	Calculate the concentration of ethanoic acid in the vinegar.	[2]
(f)	Potassium hydroxide solutions can react with carbon dioxide from the air. The solution was made one day prior to using it in the titration.	
	(i) State the type of error that would result from the student's approach.	[1]



(Question	5	continued)
(QueStion	J	continued

	(ii)	Predict, giving a reason, the effect of this error on the calculated concentration of ethanoic acid in 5(e).	[2]
	The elect		
6.	i ne elect	ron configuration of copper makes it a useful metal.	
		termine the frequency of a photon that will cause the first ionization of copper. Use stions 1, 2 and 8 of the data booklet.	[2]
	(b) Exp	plain why a copper(II) solution is blue, using section 17 of the data booklet.	[3]



Turn over

(Question 6 continued)

(c)	Copper plat	ting can be	used to im	prove the c	conductivity	of an ob	piect
()	Coppoi piai	iiig oaii bo	acca to iiii		Joinadouvity	OI GII OK	,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,

should be placed. [1]

State, giving your reason, at which electrode the object being electroplated

- **7.** Nitrogen monoxide reacts with oxygen gas to form nitrogen dioxide.
 - (a) The following experimental data was obtained.

Experiment	Initial [NO] / mol dm ⁻³	Initial [O ₂] / mol dm ⁻³	Initial rate / mol dm ⁻³ s ⁻¹
1	0.0100	0.0300	2.13×10^{-2}
2	0.0100	0.0600	4.26×10^{-2}
3	0.0300	0.0300	1.92×10^{-1}

Deduce the partial order of reaction with respect to nitrogen monoxide and oxygen. [2]

NC):
O ₂	:



(Question 7 continued)

(b) Deduce, giving a reason, whether the following mechanism is possible.

First step:	$2NO(g) \rightarrow N_2O_2(g)$	slow	
Second step:	$N_2O_2(g) + O_2(g) \rightleftharpoons 2NO_2(g)$	fast	[1]

References: NIST Mass Spectrometry Data Center Collection © 2014 copyright by the U.S. Secretary of Commerce on behalf of the United States of America. All rights reserved. COBLENTZ SOCIETY. Collection © 2018 copyright by the U.S. Secretary of Commerce on behalf of the United 2.(d) States of America. All rights reserved.

Please do not write on this page.

Answers written on this page will not be marked.



Please do not write on this page.

Answers written on this page will not be marked.

