

### Chemistry Standard level Paper 2

Thursday 12 May 2016 (morning)

	Car	ıdida	te se	ssior	num	nber		
							ļ	

1 hour 15 minutes

#### Instructions to candidates

- Write your session number in the boxes above.
- Do not open this examination paper until instructed to do so.
- · Answer all questions.
- Write your answers in the boxes provided.
- A calculator is required for this paper.
- A clean copy of the **Chemistry data dooklet** is required for this paper.
- The maximum mark for this examination paper is [50 marks].

International Baccalaureate Baccalaureate Baccalaureat International Bachillerato Internacional

Answer <b>all</b> questions.	Write your answers in the	boxes provided.
------------------------------	---------------------------	-----------------

1.	Phosphine (IUPAC name phosphane) is a hydride of phosphorus, with the formula PH <sub>3</sub> .												
	(a)	(i)	Draw a Lewis (electron dot) structure of phosphine.	[1]									
		(ii)	Outline whether you expect the bonds in phosphine to be polar or non-polar, giving a brief reason.	[1]									
		(iii)	Explain why the phosphine molecule is not planar.	[2]									
		(iv)	Phosphine has a much greater molar mass than ammonia. Explain why phosphine has a significantly lower boiling point than ammonia.	[2]									



#### (Question 1 continued)

(b) Phosphine is usually prepared by heating white phosphorus, one of the allotropes of phosphorus, with concentrated aqueous sodium hydroxide. The equation for the reaction is:

$${\rm P_4(s)} + 3{\rm OH^-(aq)} + 3{\rm H_2O\,(l)} \rightarrow {\rm PH_3(g)} + 3{\rm H_2PO_2^-(aq)}$$

(	i)	Identify	one other	element that	has allotrop	es and list two	of its allotro	pes. [2	21
١.	'''	ideritiiy		CICITICITE UTA	t Has allottop	Co and not two	or its anotio	pco. [/	<u>- 1</u>

Element:	
Allotrope 1:	
Allotrope 2:	
(ii) The first reagent is written as P <sub>4</sub> , not 4P. Describe the difference between P <sub>4</sub> and 4P.	[1]
(iii) The ion H <sub>2</sub> PO <sub>2</sub> <sup>-</sup> is amphiprotic. Outline what is meant by amphiprotic, giving the formulas of <b>both</b> species it is converted to when it behaves in this manner.	[2]



**Turn over** 

#### (Question 1 continued)

(iv)	State the oxidation state of phosphorus in $P_4$ and $H_2PO_2^-$ .	[2]
------	--	-----

P <sub>4</sub> :
H <sub>2</sub> PO <sub>2</sub> <sup>-</sup> :

(v) Oxidation is now defined in terms of change of oxidation number. Explore how earlier definitions of oxidation and reduction may have led to conflicting answers for the conversion of  $P_4$  to  $H_2PO_2^-$  and the way in which the use of oxidation numbers has resolved this.

[3]

• •		 •	•	•	 •	•	•	•	•	•	•	•	 •	•	•	•	•	 •	•	•	•	•	 •	•	•	 •	•	 •	•	 •	•	 •	•	•	 •	•	•	•	 •	•	•	 •	•		 •	
• •	•		٠	•	 •	•	•	•	•	•			 	•	٠	•	•	 •	٠	•	٠	•	 ٠	•	•	 ٠	•	 ٠	٠	 •	٠	 •	•	•		•	•	•	 ٠	٠	•	 •	٠		 ٠	
	•	 •	•	•		•	•	•	•				 		٠	٠	•	 •	٠	•	•	•	 •	•	•	 •	•	 ٠	•	 •	•	 •	•	•	 •	•	•		 •	•	•	 •	•		 ٠	۰
	•	 •	•	•		•	•	•	•				 		٠	٠	•	 •	٠	•	•	•	 •	•	•	 •	•	 ٠	•	 •	•	 •	•	•	 •	•	•		 •	•	•	 •	•		 ٠	٠
	•	 •	•	•		•	•	•	•				 		٠	٠	•	 •	•	•	•	•	 ٠	•	•	 •	•	 •	•	 •	•	 •	•	•	 •	•	•		 •	•	•	 •	•		 ٠	۰
	•	 •	•	•			•	•	•				 		٠	•	•	 •	٠	•	•	•	 ٠	•	•	 •	•	 ٠	•	 •	•	 •	•	•	 •	٠	•	•	 •	•	•	 ٠	•		 •	•
	٠		٠	•		•	٠	٠	•	•	•		 	•	٠	٠			٠	•	•	-	 ٠	•		 •	•	 ٠	٠		٠	 •	•	•	 -	٠	•	-	 •	٠		 ٠	•	-	 ٠	۰

(c) 2.478 g of white phosphorus was used to make phosphine according to the equation:

$$P_4(s) + 3OH^-(aq) + 3H_2O(l) \rightarrow PH_3(g) + 3H_2PO_2^-(aq)$$

(i) Calculate the amount, in mol, of white phosphorus used. [1]





# (Question 1 continued)

(ii) This phosphorus was reacted with 100.0 cm³ of 5.00 mol dm⁻³ aqueous sodium hydroxide. Deduce, showing your working, which was the limiting reagent.	[1]
(iii) Determine the excess amount, in mol, of the other reagent.	[1]
(iv) Determine the volume of phosphine, measured in cm³ at standard temperature and pressure, that was produced.	[1]



2.	_	ırities ( water.	cause phosphine to ignite spontaneously in air to form an oxide of phosphorus	
	(a)	(i)	200.0 g of air was heated by the energy from the complete combustion of 1.00 mol phosphine. Calculate the temperature rise using section 1 of the data booklet and the data below.	[1]
			Standard enthalpy of combustion of phosphine, $\Delta H_c^{\ominus} = -750 \text{kJ}\text{mol}^{-1}$	
			Specific heat capacity of air = $1.00 \mathrm{Jg^{-1}K^{-1}} = 1.00 \mathrm{kJkg^{-1}K^{-1}}$	
		(ii)	The oxide formed in the reaction with air contains 43.6% phosphorus by mass. Determine the empirical formula of the oxide, showing your method.	[3]
		(iii)	The molar mass of the oxide is approximately 285 g mol <sup>-1</sup> . Determine the molecular formula of the oxide.	[1]



(Question	2	continued)

(b)	(i)	State the equation for the reaction of this oxide of phosphorus with water.	[1]
	(ii)	Predict how dissolving an oxide of phosphorus would affect the pH and electrical conductivity of water.	[1]
pH:			
Elec	trical (	conductivity:	
	(iii)	Suggest why oxides of phosphorus are not major contributors to acid deposition.	[1]



Turn over

[2]

# (Question 2 continued)

(iv)	The levels of sulfur dioxide, a major contributor to acid deposition, c	an be
	minimized by either pre-combustion and post-combustion methods.	Outline one
	technique of each method.	

Pre-combustion:		
Post-combustion:		



	$\sim$	
_	ч	_

**3.** Phosgene, COCl<sub>2</sub>, is usually produced by the reaction between carbon monoxide and chlorine according to the equation:

$$CO(g) + Cl_2(g) \rightleftharpoons COCl_2(g)$$
  $\Delta H = -108 \text{ kJ}$ 

(a) (i) Deduce the equilibrium constant expression,  $K_c$ , for this reaction.

[1]


(ii) State the effect of an increase in the total pressure on the equilibrium constant,  $K_c$ .

[1]


(b) (i) Sketch the potential energy profile for the synthesis of phosgene, using the axes given, indicating both the enthalpy of reaction and activation energy.

[2]



Progress of reaction

(ii) This reaction is normally carried out using a catalyst. Draw a dotted line labelled "Catalysed" on the diagram above to indicate the effect of the catalyst.

(This question continues on the following page)




[1]

### (Question 3 continued)

(iii) Sketch and label a second Maxwell–Boltzmann energy distribution curve representing the same system but at a higher temperature,  $T_{\text{higher}}$ .

[1]




/i. /	Explain why an increase in temperature increases the rate of this reaction.	[2]
(IV	) — Explain why an increase in lemberature increases the rate of this reaction	1/1
(	= xpiam mily an moreage in temperature moreages the rate of the reaction.	[-]

		٠										 	 	•				 	•			 	•									 		



**4.** Alkenes are widely used in the production of polymers. The compound **A**, shown below, is used in the manufacture of synthetic rubber.



(a)	(i)	State the name, applying IUPAC rules, of compound <b>A</b> .	[1]
	(ii)	Draw a section, showing three repeating units, of the polymer that can be formed from compound <b>A</b> .	[1]
	(iii)	Compound <b>A</b> is flammable. Formulate the equation for its complete combustion.	[1]


(This question continues on the following page)



**Turn over** 

# (Question 4 continued)

(b) Compound **B** is related to compound **A**.



(1)	the same way as compound <b>A</b> and compound <b>B</b> .	[1]
(ii)	Suggest a chemical test to distinguish between compound <b>A</b> and compound <b>B</b> , giving the observation you would expect for each.	[2]
Test:		
Observati	on with <b>A</b> :	
Observati	on with <b>B</b> :	



[2]

[1]

[2]

#### (Question 4 continued)

(iii) Spectroscopic methods could also be used to distinguish between compounds **A** and **B**.

Predict one difference in the IR spectra **and** one difference in the <sup>1</sup>H NMR spectra of these compounds, using sections 26 and 27 of the data booklet.

<sup>1</sup>H NMR spectra:

(c) A sample of compound **A** was prepared in which the <sup>12</sup>C in the CH, group was replaced

- (c) A sample of compound  $\bf A$  was prepared in which the  $^{12}{\rm C}$  in the CH $_2$  group was replaced by  $^{13}{\rm C}$ .
  - (i) State the main difference between the mass spectrum of this sample and that of normal compound **A**.

.....

(ii) State the structure of the nucleus and the orbital diagram of <sup>13</sup>C in its ground state.

Orbital diagram

2s 2n



**Turn over** 

# (Question 4 continued)

(d) Draw a 1s atom	ic orbital and a 2p atomic orbital.	[1]
1s:	2p:	



Please **do not** write on this page.

Answers written on this page will not be marked.



Please **do not** write on this page.

Answers written on this page will not be marked.

